

NOTE XVIII.

DESCRIPTION OF A NEW SPECIES OF THE
LEPIDOPTEROUS GENUS THESTIAS

BY

R. VAN EECKE.*Thestias Weelei*, nov. spec. ♀.

Uppersurface of body black-gray; undersurface of head with palpi and of thorax pale yellow, that of abdomen white; legs yellow, tarsi brownish; antennae black above, white beneath, the tip pale brown; eyes brown.

The base of the forewing is covered with a grayish brown dust; the top is equally black-brown with a narrow short orange-yellow slightly curved band, beginning at the centre of the costa (where the colour is more pure yellow) and followed by two transverse larger orange-yellow spots, of which the first is provided near the outer margin with a black-brown spot; the second is dissolved into two small orange-yellow spots. The colour of the undescribed triangular part of the forewing is yellowish white.

The hindwings are yellowish white, with a broad fallow black-brown margin; the line of demarkation between the two colours is crenulated; the extreme base is diffusely covered with a grayish brown dust.

The undersurface of the wings is light yellow, marbled with brown exactly like in the ♀ of *Th. Piepersii* Snell. *Th. Weelei* however has in the brown spots a small whitish centre, situated upon the nervures. Moreover the new species is allied to *Th. malum sinicum* Thieme, from the island of Nias.

Wing-expansion 50 mm.

The Leyden Museum possesses one female specimen, found in the collection brought together in the Dutch East-Indies by the late Dr. H. W. van der Weele. It has been captured by him in the island of Sabang, north of Sumatra, on 21 April 1909.

Leyden Museum, February 1912.

Notes from the Leyden Museum, Vol. XXXIV.